

Cleaning the House

Use an all-purpose household cleaner. Follow the directions on the label, and:

- Wet the surface well and clean using good friction
- Allow the surface to air dry
- Pay special attention to areas that may be soiled with feces such as the toilet and sink. Remove any visible feces, and then clean as described above

Cleaning Clothes/Other Fabric

Wash clothes/fabric separately if they are heavily soiled with feces:

- Rinse off feces
- Clean in a hot water cycle with soap
- Dry items in the dryer on high heat, if possible
- Dry clean where appropriate

Cleaning dishes

Regular cleaning – use the dishwasher or clean by hand with soap and water.

Adapted from the Ottawa Hospital *C. difficile* Fact Sheet and Provincial Infectious Diseases Advisory Committee (PIDAC) Best Practice Document for the Management of *Clostridium difficile* in all Health Care Settings (January 2009)

Date of Issue: May 2014

Date of Revision:

Document #: PMH175



Clostridium difficile Infection (CDI) Client Information

What is *Clostridium difficile* (*C. difficile* or *C. diff*)?

C. difficile is just one of the many types of bacteria that can be found in the environment and the bowel. *C. difficile* is the most common cause of infectious diarrhea in hospitals and long-term care homes. It has been a known cause of health care-associated diarrhea for about 30 years.

What is *Clostridium difficile* Infection (CDI)?

For most people, *C. difficile* does not pose a health risk. When *C. difficile* bacteria grow in the bowel, it produces toxins. These toxins can damage the bowel and cause diarrhea, causing a disease known as *Clostridium difficile* infection (CDI). The effects of CDI are usually mild but sometimes can be more severe. Symptoms can range from mild or severe diarrhea to high fever, abdominal cramping, abdominal pain and dehydration. In severe cases, surgery may be needed, and in extreme cases CDI may cause death.

What causes *Clostridium difficile* Infection (CDI)?

C. difficile Infection (CDI) can sometimes occur when antibiotics are prescribed. Antibiotics work by killing off bacteria – the bad bacteria – but also good bacteria. Without the presence of the typical “good” bowel bacteria, the *C. difficile* bacteria may start to grow and produce toxins that can cause CDI.



What are the Risk Factors for CDI?

Certain people are at increased risk for acquiring CDI. These risk factors include:

- A history of antibiotic usage
- Bowel surgery
- Chemotherapy
- Prolonged hospitalization

Additional risk factors that predispose some people to develop more severe disease include:

- Increased age
- Serious underlying illness or debilitation

How Will Your Doctor Know That You Have CDI?

If you have symptoms of CDI, your doctor will ask for a sample of your watery stool. The laboratory will test the stool to see if *C. difficile* toxins are present.

How is CDI Treated?

Treatment depends on how sick you are. People with mild symptoms may not need treatment. For more severe disease, antibiotics are required. Treatment of CDI should be initiated based on the individual's risk factors and symptoms. Appropriate treatment will be determined by your doctor.

How Does CDI Spread?

When a person has CDI, the bacteria in the stool can contaminate surfaces such as toilets, handles, bedpans, or commode chairs. When touching these items our hands can become contaminated. If we then touch our mouth without washing our hands, we can become infected. Our soiled hands can also spread the bacteria to other surfaces. Washing our hands with soap and water is the best way to prevent the spread of this germ.

By always washing your hands and practicing good hygiene, you can greatly reduce your chances of picking up **any** bacteria – not just *C. difficile*.

What happens if I get CDI while I am in a health care facility?

You will be put on special precautions until you are free from diarrhea for at least two days. Your activities outside the room may be restricted. All health care staff who enter your room must wear a gown and gloves. Everyone **MUST** wash their hands when leaving your room.

Always wash your hands with soap and water after using the bathroom and before eating. As well, a thorough cleaning of your room and equipment will be done to prevent spread of the infection. Ask your visitors to check in with nursing staff if they have not done so already.

What Should I Do at Home?

Healthy people like your family and friends who are not taking antibiotics are at very low risk of getting CDI.

If you are receiving care in your home, health care workers may wear a gown and gloves depending on the situation.

It is very important that you take all your medications as prescribed by your doctor. You should not use any drugs from the drugstore that will stop your diarrhea. If diarrhea persists or comes back, contact your doctor.

Hand Hygiene

Wash your hands for at least 15 seconds with soap and water:

- After using the toilet
- After touching dirty surfaces
- Before eating
- Before preparing meals